

# Diffusion, optimal transport, and Ricci curvature

Giuseppe Savaré (*Universita' di Pavia*)

Starting from the pioneering paper of OTTO-VILLANI [11], the link between optimal transport and Ricci curvature in smooth Riemannian geometry has been deeply studied [8, 13]. Among the various functional and analytic applications, the point of view of Optimal Transport has played a crucial role in the LOTT-STURM-VILLANI [10, 12] formulation of a “synthetic” notion of lower Ricci curvature bound, which has been extended from the realm of smooth Riemannian manifolds to the general framework of metric measure spaces  $(X, \mathbf{d}, \mathbf{m})$ , i.e. (separable, complete) metric spaces endowed with a locally finite Borel measure  $\mathbf{m}$ .

Lower Ricci curvature bounds can also be captured by the celebrated BAKRY-ÉMERY [6] approach based on Markov semigroups, diffusion operators and  $\Gamma$ -calculus for local Dirichlet forms [7].

We will discuss a series of recent contributions [3, 4, 5, 9, 1] showing the link between the two approaches and the metric-variational theory of gradient flows [2] and diffusion equations. As a byproduct, when the Cheeger energy on  $(X, \mathbf{d}, \mathbf{m})$  is quadratic (or, equivalently, the Sobolev space  $W^{1,2}(X, \mathbf{d}, \mathbf{m})$  is Hilbertian), we will show that the two approaches lead to essentially equivalent definitions and to a nice geometric framework suitable for deep analytic results.

## References

- [1] L. Ambrosio, A. Mondino, and G. Savaré. Nonlinear diffusion equations and curvature conditions in metric measure spaces. *ArXiv* 1509.07273, 2015.
- [2] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, second edition, 2008.
- [3] L. Ambrosio, N. Gigli, and G. Savaré. Calculus and heat flow in metric measure spaces and applications to spaces with ricci bounds from below. *Invent. Math.*, 195(2):289–391, 2014.
- [4] L. Ambrosio, N. Gigli, and G. Savaré. Metric measure spaces with Riemannian Ricci curvature bounded from below. *Duke Math. J.*, 163(7):1405–1490, 2014.
- [5] L. Ambrosio, N. Gigli, and G. Savaré. Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds. *Ann. Probab.*, 43(1):339–404, 2015.

- [6] D. Bakry and M. Émery. Diffusions hypercontractives. In *Séminaire de probabilités, XIX, 1983/84*, volume 1123, pages 177–206. Springer, Berlin, 1985.
- [7] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften*. Springer, Cham, 2014.
- [8] D. Cordero-Erausquin, R.J. McCann, and M. Schmuckenschläger. A Riemannian interpolation inequality à la Borell, Brascamp and Lieb. *Invent. Math.*, 146(2):219–257, 2001.
- [9] M. Erbar, K. Kuwada, and K.-T. Sturm. On the Equivalence of the Entropic Curvature-Dimension Condition and Bochner’s Inequality on Metric Measure Spaces. *ArXiv* 1303.4382, 2013. To appear on *Invent. Math.*
- [10] J. Lott and C. Villani. Ricci curvature for metric-measure spaces via optimal transport. *Ann. of Math. (2)*, 169(3):903–991, 2009.
- [11] F. Otto and C. Villani. Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *J. Funct. Anal.*, 173(2):361–400, 2000.
- [12] K.T. Sturm. On the geometry of metric measure spaces. I, II. *Acta Math.*, 196(1):65–131 and 133–177, 2006.
- [13] K.T. Sturm and M.K. von Renesse. Transport inequalities, gradient estimates, entropy, and Ricci curvature. *Comm. Pure Appl. Math.*, 58(7):923–940, 2005.